

HDP Session 1

1. Concentration: Eg. X_i iid X , $\mathbb{E}X = \mu$, $\text{Var}(X) < \infty$

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{p} \mathbb{E}X = \mathbb{E}\bar{X}_n = \mu$$

Asymptotic Frequentist Approach to Prob.

Q: At what rate?

- Markov's Ineq: $X \geq 0$ a.s., then $\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}X}{a}$, $\mathbb{E}X < \infty$, $\forall a > 0$
- Chebychev's Ineq: $\mathbb{E}X^2 < \infty$, then $\mathbb{P}(|X - \mathbb{E}X| \geq a) \leq \frac{\text{Var}(X)}{a^2}$, $\forall a > 0$

$$\textcircled{*} \rightarrow \checkmark \quad \mathbb{P}(|\bar{X}_n - \mathbb{E}\bar{X}_n| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X}_n)}{\varepsilon^2} = \frac{\text{Var}(X)}{n\varepsilon^2} \sim O\left(\frac{1}{n}\right) \xrightarrow{\text{Non-asymptotic.}} e^{-n\varepsilon^2}$$

1) Can the rate of conv. be improved? — For some special classes of rvs.

Subgaussian

Subexponential

$$\bar{X}_n \xrightarrow{p} \mathbb{E}\bar{X}_n \quad ; \quad \bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

2) For what other function $f_n: \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies

$$\textcircled{*} \text{ Conc. ineq. } \rightarrow \mathbb{P}(|f_n(X_1, \dots, X_n) - \mathbb{E}f_n(X_1, \dots, X_n)| \geq \varepsilon) \leq \frac{\text{Var}(f_n(X_1, \dots, X_n))}{\varepsilon^2}$$

$$\text{eg. } f_n(X_1, \dots, X_n) = \frac{X_1^2 + \dots + X_n^2}{n}, \quad X_i \text{ iid } N(0,1)$$
$$f_n(X_1, \dots, X_n) \xrightarrow{p} \mathbb{E}(f_n(X_1, \dots, X_n))$$

→ Martingale methods

Indeed, showing conc. is sometimes the easy part.

$\mathbb{E}f(X_1, \dots, X_n) \rightarrow$ harder. ←

✓ 2. Supremum: $\{X_t \mid t \in T\} \rightarrow$ Some collection of rvs.

$$\mathbb{P} \left(\sup_{t \in T} X_t \leq x \right), \quad \mathbb{E} \sup_{t \in T} X_t$$

⊛ Exc: $X_1, X_2, \dots$ iid $N(0,1)$. Estimate $\mathbb{E} \left(\sup_{i=1}^n X_i \right)$.

Thm. (Fisher - Tippet - Gnedenko) X_i iid $\underline{X}$ with some good properties.

/ Then $\exists$ sequences $\{a_n\}, \{b_n\}$ s.t.

Extreme
Value
Theory.

$$\frac{\max_{i=1}^n X_i - a_n}{b_n} \xrightarrow{d} \zeta$$

Gumbel distribution

$$\cdot F_G(x) = e^{-e^{-x}}$$

$$Y_n \xrightarrow{d} Y$$

$$F_{Y_n} \rightarrow F_Y \text{ ptwise at all cont. pts of } F_Y.$$

What is the non-asymptotic version of this?

3) Universality: $\cdot$ CLT: X_i iid X , $\mathbb{E}X = \mu$, $\text{Var} X = \sigma^2$

$$\sqrt{n}(\bar{X}_n - \mu) \xrightarrow{d} N(0, \sigma^2)$$

→ Fourier Analytic Method

→ Method of Moments $\rightsquigarrow$ Lindeberg Trick $\star$

2. Moment Methods - FMM & SMM

Events $\xleftrightarrow{?}$ Random Variable $;$ Ω

$$A \in \Omega \rightsquigarrow \mathbb{1}_A(\omega) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{o.w.} \end{cases}$$

Defⁿ: (Erdős - Renyi RGs): $G(n, p)$ is a graph of n vertices $\{1, 2, \dots, n\} = [n]$, s.t.

$\{i, j\} \in E(G(n, p))$ w.p. p indep of all other edges

$G(3, p)$ 

eg. Consider $G(n, p)$, and let $A_i = \{i^{\text{th}} \text{ vertex is isolated}\}$.

$$\text{Let } Y_{i,j} = 1_{\{\{i,j\} \in E(G(n,p))\}}$$

$$\begin{aligned} \underline{P(A_i)} &= P(Y_{i,1}, Y_{i,2}, \dots, Y_{i,n} = 0) \\ &= P(Y_{i,1} + \dots + Y_{i,n} = 0) \rightsquigarrow \text{Will be expressed in terms of a conc. ineq.} \\ \mathbb{E} Y_{i,1} = p &\rightarrow \leq P\left(\left|\sum_{j \neq n} Y_{i,j} - (n-1)p\right| \geq (n-1)p\right) \\ &\leq \dots \end{aligned}$$

⊛ First Moment method: $X \geq 0$ a.s. and $\mathbb{E} X < \infty$, X - integer valued

Then

$$\begin{aligned} \mathbb{E} X &= \int_0^\infty P(X \geq t) dt = \int_0^a P(X \geq t) dt + \int_a^\infty P(X \geq t) dt \\ &\geq \int_0^a P(X \geq t) dt \geq a \cdot P(X \geq a) \end{aligned}$$

$$\therefore P(X \geq a) \leq \frac{\mathbb{E} X}{a} \quad [\text{Markov's Ineq.}]$$

$$\boxed{P(X > 0) = P(X \geq 1) \leq \mathbb{E} X} \rightarrow \textcircled{I}$$

⊛ Second Moment Method: $X \geq 0$ a.s. and $\mathbb{E} X^2 < \infty$, then

$$\begin{aligned} (\mathbb{E} X)^2 &= (\mathbb{E}(X 1_{X>0}))^2 \leq (\mathbb{E} X^2) \cdot \mathbb{E}(1_{X>0}) \\ &= \mathbb{E} X^2 \cdot P(X > 0) \end{aligned}$$

$$P(X > 0) \geq \frac{(\mathbb{E} X)^2}{\mathbb{E} X^2}$$

★ $0 \rightarrow \frac{(\mathbb{E} X)^2}{\mathbb{E} X^2} \leq P(X > 0) \leq \mathbb{E} X$ for $X - \mathbb{N}_0$ valued rv.

$$\begin{aligned} \mathbb{N} &= \{1, 2, \dots\} \\ \mathbb{N}_0 &= \{0, 1, 2, \dots\} \end{aligned}$$


Ex 1.: $G(n, p)$: $\mathbb{P}(A_i)$, $A_i = i^{\text{th}}$ vertex is isolated

$$\rightarrow \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \mathbb{P}(G(n, p) \text{ has an isolated vertex})$$

$$= \underline{\underline{\mathbb{P}(N > 0)}}$$

• $N = \#$ isolated vertices.

$$N = \sum_{i=1}^n N_i, \quad N_i = \mathbb{1}[i^{\text{th}} \text{ vertex is isolated}]$$



$$N_i \sim \text{Ber}((1-p)^{n-1}) \quad \forall i$$

$$\mathbb{E}N = \sum_{i=1}^n \mathbb{E}N_i = n(1-p)^{n-1}$$

$$\leq e^{\log n} \cdot e^{-p(n-1)}$$

$$= \exp\left(\underbrace{\log n - (n-1)p}_{\text{negative for large } n}\right)$$

$$(1-x) \leq e^{-x} \quad \forall x \in \mathbb{R}$$

If $p \geq c \frac{\log n}{n}$ for some $c > 1$, then $\mathbb{E}N \rightarrow 0$

$$\mathbb{P}(N > 0) \leq \mathbb{E}N \rightarrow 0$$

•• For $p > c \frac{\log n}{n}$ for some $c > 0$, we have $\mathbb{P}(G(n, p) \text{ isol vert}) = o(1)$.

• 2nd moment method: N_i 's are not independent

$$N_i = \mathbb{1}[i \text{ is isolated}]$$

$$\mathbb{E}(N^2) = \mathbb{E}\left(\left(\sum_{i=1}^n N_i\right)\left(\sum_{j=1}^n N_j\right)\right)$$


$$= \mathbb{E}\left(\sum_{i=1}^n N_i^2\right) + 2 \mathbb{E}\left(\sum_{i < j} N_i N_j\right)$$

$$= \underbrace{\sum_{i=1}^n \mathbb{E}N_i}_{n(1-p)^{n-1}} + 2 \underbrace{\sum_{i < j} \mathbb{E}(N_i N_j)}_{n(n-1)(1-p)^{2n-3}}$$

$$\mathbb{E}(N_i N_j) = \mathbb{E}(\mathbb{1}[i \text{ is isolated}] \cdot \mathbb{1}[j \text{ is isolated}])$$

$$= \mathbb{E}(\mathbb{1}[i, j \text{ are isolated}])$$

$$= (1-p)^{2n-3}$$



$$P(N > 0) \geq \frac{(\mathbb{E}N)^2}{\mathbb{E}(N^2)} = \frac{(n(1-p)^{n-1})^2}{n(n-1)(1-p)^{2n-3} + (n(1-p)^{n-1})}$$

$$\geq \frac{n^2 (1-p)^{2n-2}}{n^2 (1-p)^{2n-3} + n(1-p)^{n-1}}$$

$$\frac{1}{P(N > 0)} \leq \frac{1}{1-p} + \frac{1}{n(1-p)^{n-1}} \quad ; \rightarrow \textcircled{2}$$

★ Result: $n(1-p)^{n-1} \rightarrow \begin{cases} 0 & \text{if } p > C \frac{\log n}{n} \text{ for } C > 1 \\ \infty & \text{if } p < c \frac{\log n}{n} \text{ for } c < 1 \end{cases}$

∴ For $p < c \frac{\log n}{n}$, we have

$$\frac{1}{P(N > 0)} \leq 1 + o(1) \Rightarrow P(N > 0) \geq 1 - o(1)$$


$$\therefore \lim_{n \rightarrow \infty} P(N > 0) = 1 \quad \square$$

☆ We have show that

$\{\exists \text{ an isolated vertex in } G(n, p)\}$

undergoes a phase transition at $p = \frac{\log n}{n}$

Phase
Transition



Eg 2. $P(G(n, p) \text{ is connected}) \geq P(G(n, p) \text{ has no isol. vertices})$

∴ If $p \ll \frac{\log n}{n}$, then $P(G(n, p) \text{ is connected}) = o(1)$

$p \gg \frac{\log n}{n}$, then $P(\underline{G(n, p)} \text{ is conn.}) = 1 - o(1)$.

Let Y_k be the # of connected components of size k , $k=1, 2, \dots, n/2$.
Then,

$$\mathbb{P}(G(n,p) \text{ is not conn.}) = \mathbb{P}\left(\underbrace{\sum_{k=1}^{n/2} Y_k}_{>0}\right) \leq \sum_{k=1}^{n/2} \mathbb{E} Y_k.$$

$$\bullet \quad \mathbb{E} Y_k \leq \binom{n}{k} (1-p)^{k(n-k)}.$$

$$\stackrel{||}{=} \mathbb{E} \left(\sum_{A \in \binom{[n]}{k}} Y_k^A \right)$$

subsets of n of size k

$$Y_k^A = \mathbb{1}[G_A \text{ is a conn. comp}]$$



$$\circ \circ \quad \mathbb{P}(G(n,p) \text{ is disconn.}) \leq \sum_{k=1}^{n/2} \binom{n}{k} (1-p)^{k(n-k)}$$

$$\leq \sum_{k=1}^{n/2} n^k (1-p)^{k(n-k)}$$


$$k \leq \frac{n}{2} \Rightarrow n-k \geq \frac{n}{2}$$

$$\leq \sum_{k=1}^{n/2} n^k \left((1-p)^{n/2} \right)^k \rightarrow 0$$

as $n \rightarrow \infty$ for $p > C \frac{\log n}{n}$

$$1-p \leq e^{-p}$$

$$\circ \circ \quad \mathbb{P}(G(n,p) \text{ is conn.}) = 1 - \underbrace{o(1)}_{\rightarrow 0} \text{ for } \underline{p > C \frac{\log n}{n}}.$$



- $\mathbb{P}(G(n,p) \text{ has no isol. vertices})$
- $\mathbb{P}(G(n,p) \text{ is connected})$

As $n \uparrow \infty$, these prob. become indistinguishable

[Sebastian Roch: Example 2...]

$$\bullet \quad |\text{SuperCr}|, |\text{SubCr}| \gg |\text{Crit}|$$

Limit of Graph (V, E)

⊛ Thm:

$$\underbrace{p = \frac{\log n + s}{n}}_{\text{, then}} \begin{cases} \text{IP}(\# \text{ isol vertices}) = \frac{1}{1+e^s} + o(1) \\ \text{IP}\left(\sum_{k=2}^{n/2} Y_k > 0\right) = \underline{o(1)} \leftarrow \end{cases}$$

